

治療計画書

療法名

アービタックス+RT

療法

1サイクル 1 週

疾患名 頭頸部がん(7~8サイクル)

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																																	
			用量(/日)	用法		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28					
		照射部位	照射量																																			
注射	1	デキサート	6.6 mg	点滴	30分																																	
		ボラミン	5 mg			●																																
		生理食塩液	100 ml																																			
注射	2	生理食塩液	50 ml	点滴	15分	●																																
注射	3	アービタックス<<初回>>	400 mg/m ²	点滴	2時間																																	
		生理食塩液	500 ml	全量500mlに調製		●																																
		アービタックス<<2回目以降>>	250 mg/m ²	点滴	1時間																																	
		生理食塩液	250 ml	全量250mlに調製		●																																
注射	4	生理食塩液<<初回>>	50 ml	点滴	1時間																																	
				1時間の状態観察		●																																
		生理食塩液<<2回目以降>>	50 ml	点滴	15分																																	
				1時間の状態観察		●																																
注射	5																																					
注射	6																																					
注射	7																																					
注射	8																																					
注射	9																																					
注射	10																																					
内服	1																																					
内服	2																																					
内服	3																																					
内服	4																																					
内服	5																																					
内服	6																																					
放射線		2サイクル目より併用	60-76.8 Gy	1日1-2回		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	

特記事項

アービタックス

- ・至適濃度: 0.83mg/ml~4mg/ml
- ・治療開始前、治療中、終了後に電解質(Mg、K、Ca)のモニタリングを行う。
- ・皮膚障害対策は「適正使用ガイド」を参照
- ・infusion reactionを発現することがあるため、アービタックス投与中及び投与終了後は少なくとも1時間は観察期間(バイタルサインをモニターするなど)を設けること。
- ・2サイクル目より放射線を併用

減量規定

薬品名	アービタックス			
標準投与量	250 mg/m ²			
1段階減量	200 mg/m ²			
2段階減量	150 mg/m ²			
3段階減量	中止			