

治療計画書

療法名

アムルビシン

(ID 0) ☆7

療法

1サイクル 3~4週

疾患名 **小細胞肺癌(※4~6サイクル)**

照射線	商品名	用量(/回)	投与ルート	投与時間	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
内服		用量(/日)	用法																															
照射線	照射部位	照射量																																
注射	1	グラニセトロン デキサート	3 mg 6.6 mg	点滴	30分	●	●	●																										
注射	2	5%糖液	500 ml	点滴	2時間	●	●	●																										
注射	3	カルセド 生理食塩液	40 mg/m ² 50 ml	点滴(側管)	5分 (全開)	●	●	●																										
注射	4																																	
注射	5																																	
注射	6																																	
注射	7																																	
注射	8																																	
注射	9																																	
注射	10																																	
内服	1																																	
内服	2																																	
内服	3																																	
内服	4																																	
内服	5																																	
内服	6																																	
内服	7																																	
内服	8																																	
照射線																																		

特記事項

・副作用として血液毒性(白血球減少)が高頻度に現れる
 ・血管外漏出時には水胞性皮膚壊死を起こす
 ・次クール治療時、減量の目安としては以下を参照すること
 ①白血球数<1000、好中球数<500が4日以上継続した場合②血小板の最低値<5万の場合③発熱を伴う好中球減少(<1000)が認められた場合
 ※効果が認められる場合は、総サイクル数を超えて繰り返し投与可能

減量規定

薬品名	カルセド			
標準投与量	40 mg/m ²			
1段階減量	35 mg/m ²			
2段階減量				
3段階減量				