療法

1サイクル __3 __週

_{疾患名} 非小細胞肺がん(6サイクル)

大			7 1 7 7 7																								
注射			用量(/回)	投与ルート	投与時間	1				П	2		П			П	3					Τ.	4	Т			П
内服		商品名	用量(/日)	用法	IX T FI IFI		2	2 4	5	6			10	11 1	2 12	14		16 1	7 10	10	20			2 24	25	26 2	7 20
放射線			照射量	用冱		Ľ	2	3 4	3	0	/ 0	9	10	1111	2 13	14	15	10 1	/ 10	3 19	20	21 2	2 2	3 24	25	20 2	/ 20
放射線		照射部位		F >±	00/\		-			Н			Н			H	_	_	+	\blacksquare	\dashv	_	+	+	₩	_	+
		アロキシ	0.75 mg	点滴	30分	4																					
注	1	ファモチジン	20 mg																								
射	ľ	デキサート	19.8 mg																								
		生理食塩液	100 ml																								
		生理食塩液	50 ml	点滴	15分					П						П					П		Т				П
注				,,,,,,																							
注射	2																										
ניג						-																					
	_	° 5 1 1 5 5 1	200 / 2	F 144	0.04.88					Н						Н			+		\dashv		+	4			+
		パクリタキセル	200 mg/m²	点滴	3時間																						
注射	3	生理食塩液	500 ml			•																					
射	ľ																										
		生理食塩液	50 ml	点滴	15分																П						
注						1_																					
注射	4					•																					
-33																П											
	\vdash	カルボプラチン	XAUC6	点滴	1時間30分		+	+		H	+		H	+		H	\dagger	+	۲		\dashv	+	+	+	\forall	\top	\forall
:+		生理食塩液	250 ml	小小门	, [4]00/]											Н											
注射	5	工任及塩/仪	※最大800mg																								
ניל			※取入ouomg			-																					
		上四人上法	50 1	F '*	457					Н			Н			Ш	_	_	+		\dashv		+	4	-	_	ш
		生理食塩液	50 ml	点滴	15分																						
注射	6																										
射	1																										
																					Ш						
注射	_																										
射	<i>'</i>																										
								+		Ħ			П	_		П	7	+	T		\dashv		T	T			H
注						1																					
注射	8					-																					
ניג						-																					
							\dashv	+		Н			Н	+		Н	+	+	+		\dashv	-	+	╄	-	+	+
						-																					
注射	9																										
射																											
										Ш						Ш											Ш
注	10																										
注射	10																										
内服	1	レスタミン	50 mg	パクリタキセルの	30分前まで					H			H			H			\dagger	П	T		+	+			\forall
		デカドロン	8 mg	1日1(-2)回朝				•))	H			Н			H	+		+		\dashv		+	+			+
内服			Vg		<u> </u>		Ť			H			H			H	+	\top	+		\dashv		+	+		+	+
内服							+	+	+	H			Н	+		Н	+	+	+		\dashv	-	+	+		+	+
内服	_						+			Н			Н			Н		+	+		\dashv		+	┿	\vdash	+	+
							\dashv			\vdash			H			H	+		-		\dashv		+	+	\vdash	-	+
内服							\dashv			\vdash			H	+		\vdash	4	-	-		\dashv	4	+	+	H	-	+
内服							\vdash	_		$\vdash \vdash$			\vdash	_		Н	4	_			\dashv	_	4	4	\blacksquare	-	\sqcup
内服							\sqcup			Н			Ш			Ш	4		_		\dashv		4	4	\sqcup	1	\sqcup
内服										Ц			Ш			Ш	4	\perp	L		Щ		4	4	Ш		Ш
内服										Ц			Ш			Ш					Ш		\perp	1	Ш		Ш
放射線			Gy							Ш			Ш			Ш							Ш				
					特記事項																						╝
																											Ī
<u></u>	_				減量規定																						
薬	品	名 カルボプラチン	パクリタ=																								
標準	投	与量 AUC6	200 mg	g/ m i																							
1段	階	減量 AUC5	150 mg	g/ m i๋																							
2段	階	減量 AUC4	100 mg	g/mt																							
3段	階	減量中止	中山	E																							