

治療計画書

療法名

シスプラチン+ドセタキセル

療法

1サイクル 3~4 週

疾患名 非小細胞肺がん(4サイクル)

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
			用量(/日)	用法																													
		照射部位	照射量																														
注射	1	アロキシ	0.75 mg	点滴	30分																												
		デキサート	9.9 mg			●																											
		生理食塩液	100 ml																														
注射	2	生理食塩液	500 ml	点滴	2時間																												
		硫酸Mg補正液	8 mEq			●																											
注射	3	YDソリター-T3	500 ml	点滴	2時間	●	●	●																									
						●	●	●																									
注射	4	ドセタキセル	60 mg/m ²	点滴	1時間																												
		生理食塩液	250 ml			●																											
注射	5	生理食塩液	100 ml	点滴	30分																												
						●																											
注射	6	シスプラチン	80 mg/m ²	点滴	2時間																												
		生理食塩液	500 ml	全量500mlに調製		●																											
注射	7	YDソリター-T3	500 ml	点滴	2時間																												
		フロセミド	20 mg			●																											
注射	8	ソリューゲンF	500 ml	点滴	2時間	●	●	●																									
注射	9																																
注射	10																																
内服	1	イメンド	125 mg	抗癌剤の1時間-1時間半前		●																											
内服	2	イメンド	80 mg	1日1回朝食後		●	●																										
内服	3	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●	●	●																								
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

G3以上の神経障害が現れた場合は投与を中止する

減量規定

薬品名	シスプラチン		ドセタキセル	
標準投与量		80 mg/m ²		60 mg/m ²
1段階減量	白血球減少G4が3日以上持続、 好中球減少G4が3日以上持続、 血小板減少G4、G2の神経障害、 G3-4の肝毒性、G3の口内炎・腎障害	60 mg/m ²	白血球減少G4が3日以上持続、 好中球減少G4が3日以上持続、 血小板減少G4、G2の神経障害、 G3-4の肝毒性	45 mg/m ²