

# 治療計画書

療法名

## ドセタキセル+サイラムザ

療法

1サイクル 3 週

疾患名 切除不能な進行・再発の非小細胞肺癌

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																															
			用量(/日)	用法		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
		照射部位	照射量																																	
注射	1	デキササート	6.6 mg	点滴	30分																															
		ボラミン	5 mg			●																														
		生理食塩液	100 ml																																	
注射	2	生理食塩液	50 ml	点滴	15分																															
						●																														
注射	3	サイラムザ	10 mg/kg	点滴	※1時間																															
		生理食塩液	250 ml			●																														
注射	4	生理食塩液	100 ml	点滴	15分																															
						●																														
注射	5	ドセタキセル	60 mg/m <sup>2</sup>	点滴	1時間																															
		生理食塩液	250 ml			●																														
注射	6	生理食塩液	50 ml	点滴	15分																															
						●																														
注射	7																																			
注射	8																																			
注射	9																																			
注射	10																																			
内服	1																																			
内服	2																																			
内服	3																																			
内服	4																																			
内服	5																																			
内服	6																																			
内服	7																																			
内服	8																																			
内服	9																																			
内服	10																																			
放射線			Gy																																	

### 特記事項

※G1または2のinfusion reactionが発現した場合は2時間になるよう減速し、その後の投与においても減速した速度で投与する。

(infusion reactionが発現した場合の対応については添付文書参照)

※※詳細は添付文書参照

### 減量規定

薬品名	ドセタキセル	サイラムザ※※			
標準投与量	60 mg/m <sup>2</sup>	10 mg/kg			
1段階減量	50 mg/m <sup>2</sup>	8(6) mg/kg			
2段階減量	中止	6(5) mg/kg			
3段階減量		中止			