

治療計画書

療法名 カルボプラチン+パクリタキセル+キイトルーダ 療法 1サイクル 3 週

疾患名 切除不能な進行・再発の扁平上皮非小細胞肺癌(4サイクル)

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
			用量(/日)	用法																													
		照射部位	照射量																														
注射	1	生理食塩液	50 ml	点滴	15分	●																											
注射	2	キイトルーダ	200 mg/body	点滴	30分	●																											
		生理食塩液	100 ml																														
注射	3	生理食塩液	50 ml	点滴	15分	●																											
注射	4	アロキシ	0.75 mg	点滴	30分	●																											
		ファモチジン	20 mg																														
		デキサート	19.8 mg																														
		生理食塩液	100 ml																														
注射	5	生理食塩液	50 ml	点滴	15分	●																											
注射	6	パクリタキセル	200 mg/m ²	点滴	3時間	●																											
		生理食塩液	500 ml																														
注射	7	生理食塩液	50 ml	点滴	15分	●																											
注射	8	カルボプラチン	※AUC6	点滴	1時間30分	●																											
		生理食塩液	250 ml																														
			※最大800mg																														
注射	9	生理食塩液	50 ml	点滴	15分	●																											
注射	10																																
内服	1	レスタミン	50 mg	パクリタキセルの30分前まで		●																											
内服	2	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●●●																											
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

インラインフィルター(0.22μm以下)を使用すること。

減量規定

薬品名	カルボプラチン	パクリタキセル	キイトルーダ
標準投与量	AUC6	200 mg/m ²	200 mg
1段階減量	AUC5	150 mg/m ²	減量不可
2段階減量	AUC4	100 mg/m ²	
3段階減量	中止	中止	