

治療計画書

療法名

XELOX

療法

1サイクル

3 週

疾患名 治癒切除不能な進行・再発の胃がん

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																														
			用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
		照射部位	照射量																																
注射	1	グラニセトロン	3 mg	点滴	30分																														
		デキサート	6.6 mg			●																													
注射	2	エルプラット	130 mg/m ²	点滴	2時間																														
		5%糖液	250 ml			●																													
注射	3	生理食塩液	50 ml	点滴	15分																														
						●																													
注射	4																																		
注射	5																																		
注射	6																																		
注射	7																																		
注射	8																																		
内服	1	ゼローダ※	2000 mg/m ²	1日2回朝夕食後		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
内服	2	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●																												
内服	3																																		
内服	4																																		
内服	5																																		
内服	6																																		
内服	7																																		
内服	8																																		
内服	9																																		
内服	10																																		

特記事項

エルプラット総投与量が600mg/m²を超えた場合、エルプラットは担当医の判断で適宜休止して次のサイクルを開始する。
※ゼローダは1日目の夕食後より服用開始。

ゼローダ減量基準	1日投与量		
	初回投与量	減量段階1	減量段階2
1.36m ² 未満	2400mg/日	1800mg/日	1200mg/日
1.36m ² 以上1.41m ² 未満	3000mg/日	2400mg/日	
1.41m ² 以上1.51m ² 未満		3600mg/日	1800mg/日
1.51m ² 以上1.66m ² 未満	3000mg/日		
1.66m ² 以上1.81m ² 未満		4200mg/日	2400mg/日
1.81m ² 以上1.96m ² 未満			
1.96m ² 以上2.11m ² 未満			
2.11m ² 以上			

減量規定

薬品名	エルプラット	ゼローダ		
標準投与量	130 mg/m ²	上記参照		
1段階減量	100 mg/m ²			
2段階減量	85 mg/m ²			
3段階減量		中止(CCr≤30)		