

疾患名 腎盂がん、尿管がん、膀胱がん、尿道がん

注射 内服 放射線	Rp	商品名 照射部位	用量(回)	投与ルート	投与時間	1							2							3							4									
			用量(日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28				
注射	1	ソリューゲンF	500 ml	点滴	2時間	●	●												●																	
注射	2	メソトレキセート 生理食塩液	30 mg/m <sup>2</sup> 100 ml	点滴	30分	●													●																	●
注射	3	ソリューゲンF	500 ml	点滴	2時間	●	●																													
注射	4	アロキシ デキサート 生理食塩液	0.75 mg 13.2 mg 50 ml	点滴	30分				●																											
注射	5	生理食塩液	50 ml	点滴	15分				●																											●
注射	6	エクザール 生理食塩液	3 mg/m <sup>2</sup> 100 ml	点滴	30分				●																											●
注射	7	ドキシソルピシン 生理食塩液	30 mg/m <sup>2</sup> 100 ml	点滴	30分				●																											
注射	8	シスプラチン 生理食塩液	70 mg/m <sup>2</sup> 500 ml	点滴	2時間				●																											
注射	9	YDソリター-T1 フロセミド	500 ml 10 mg	点滴	2時間				●																											
注射	10	ソリューゲンF	500 ml	点滴	2時間				●																											
注射	11	生理食塩液	50 ml	点滴	15分																															●
内服	1	デカドロン	8 mg	1日1(-2)回朝(昼)食後				●	●	●	●																									
内服	2																																			
内服	3																																			
内服	4																																			
内服	5																																			
内服	6																																			
放射線			Gy																																	

特記事項

ドキシソルピシンは総投与量500mg/m<sup>2</sup>までとする

減量規定

薬品名	メソトレキセート	エクザール	ドキシソルピシン	シスプラチン
標準投与量	30 mg/m <sup>2</sup>	3 mg/m <sup>2</sup>	30 mg/m <sup>2</sup>	70 mg/m <sup>2</sup>
1段階減量	25 mg/m <sup>2</sup>	2.5 mg/m <sup>2</sup>	25 mg/m <sup>2</sup>	60 mg/m <sup>2</sup>
2段階減量	20 mg/m <sup>2</sup>	2 mg/m <sup>2</sup>	20 mg/m <sup>2</sup>	50 mg/m <sup>2</sup>
3段階減量	15 mg/m <sup>2</sup>	1.5 mg/m <sup>2</sup>	15 mg/m <sup>2</sup>	40 mg/m <sup>2</sup>