

治療計画書

療法名

DP(ドセタキセル+シスプラチン)

療法

1サイクル 3~4 週

疾患名 卵巣がん(3~5サイクル)

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
			用量(/日)	用法																													
		照射部位	照射量																														
注射	1	ソリューゲンF	500 ml	点滴	2時間	●																											
注射	2	ソリューゲンF	500 ml	点滴	2時間	●																											
注射	3	グラニセトロン デキサート	3 mg 9.9 mg	点滴	30分	●																											
注射	4	生理食塩液	100 ml	点滴	30分	●																											
注射	5	ドセタキセル 生理食塩液	60-75 mg/m ² 250 ml	点滴	1時間以上	●																											
注射	6	生理食塩液	50 ml	点滴	10分	●																											
注射	7	シスプラチン 生理食塩液	70 mg/m ² 500 ml	点滴 全量500mlに調製	2時間以上	●																											
注射	8	ソリューゲンG フロセミド	500 ml 10 mg	点滴	2時間	●																											
注射	9	YDソリター-T3	500 ml	点滴	2時間	●																											
注射	10																																
内服	1	イメンド	125 mg	抗癌剤の1時間-1時間半前		●																											
内服	2	イメンド	80 mg	1日1回朝食後		●●																											
内服	3	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●●●●																											
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

アレルギー 予防のため適宜デキサートの増量、ファモチジンの追加を行う。制吐剤は適宜変更。

減量規定

薬品名	標準投与量	1段階減量	2段階減量	3段階減量