

# 治療計画書

療法名 TP(パクリタキセル+シスプラチン)

療法 1サイクル 3~4週

疾患名 卵巣がん (3~5サイクル)

注射 内服 放射線	Rp	商品名	用量(回)	投与ルート	投与時間																															
			用量(日)	用法		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
		照射部位	照射量																																	
注射		ソリューゲンF	500 ml	点滴	2時間	●																														
注射	1																																			
注射	2	グラニセトロン デキサート ファモチジン	3 mg 19.8 mg 20 mg	点滴	30分	●																														
注射	3	生理食塩液	100 ml	点滴	30分	●																														
注射	4	パクリタキセル 生理食塩液	180 mg/m <sup>2</sup> 500 ml	点滴	3時間	●																														
注射	5	生理食塩液	50 ml	点滴	10分	●																														
注射	6	シスプラチン 生理食塩液	70 mg/m <sup>2</sup> 500 ml	点滴	2時間以上	●																														
注射	7	ソリューゲンG フロセミド	500 ml 10 mg	点滴	2時間	●																														
注射	8	YDソリターT3	500 ml	点滴	2時間	●																														
注射	9																																			
注射	10																																			
内服	1	レスタミン	50 mg	パクリタキセルの30分前まで		●																														
内服	2	イメンド	125 mg	抗癌剤の1時間-1時間半前		●																														
内服	3	イメンド	80 mg	1日1回朝食後		●	●																													
内服	4	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●	●	●																											
内服	5																																			
内服	6																																			
内服	7																																			
内服	8																																			
内服	9																																			
内服	10																																			
放射線			Gy																																	

特記事項

制吐剤は適宜変更。

減量規定

薬品名	シスプラチン		
標準投与量	100 %		
1段階減量	45 < CCr ≤ 60	75 %	
2段階減量	30 < CCr ≤ 45	50 %	
3段階減量	CCr ≤ 30	中止	