

治療計画書

療法名 JALSG AML-201 強力寛解後 第1コース(Ara-C+M)

療法

1サイクル _____ 日

疾患名 急性骨髄性白血病

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																															
			用量(/日)	用法			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
		照射部位	照射量																																	
注射	1	生理食塩液	200 ml	点滴	キープ	●	●	●	●	●																										
注射	2	グラニセトロン	3 mg	点滴(側管)	30分	●	●	●																												
		デキサート	6.6 mg																																	
注射	3	デキサート	6.6 mg	点滴(側管)	15分																															
		生理食塩液	50 ml						●	●																										
注射	4	ノバントロン	7 mg/m ²	点滴(側管)	30分	●	●	●																												
		生理食塩液	100 ml																																	
注射	5	キロサイド	200 mg/m ²	点滴(側管)	24時間	●	●	●	●	●																										
		生理食塩液	500 ml																																	
注射	6	生理食塩液	50 ml	点滴(側管)	ルートフラッシュ																															
注射	7																																			
注射	8																																			
注射	9																																			
注射	10																																			
内服	1																																			
内服	2																																			
内服	3																																			
内服	4																																			
内服	5																																			
内服	6																																			
内服	7																																			
内服	8																																			
内服	9																																			
内服	10																																			
放射線			Gy																																	

特記事項

前治療の影響がなくなり、十分に骨髄機能が回復した後に施行

減量規定

薬品名	標準投与量	1段階減量	2段階減量	3段階減量