

治療計画書

療法名 JALSG AML-201(高齢者)寛解後 第1コース(BHAC+M) 療法 1サイクル 日

疾患名 急性骨髄性白血病

照射線	Rp	商品名	用量(/回)	投与ルート	投与時間	1							2							3							4						
			用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
照射線	1	照射部位	照射量																														
		生理食塩液	500 ml	点滴	キープ	●	●	●	●	●																							
注射	2	商品名	用量	投与ルート	投与時間																												
		グラニセトロン	3 mg	点滴(側管)	30分	●	●	●	●	●																							
注射	3	商品名	用量	投与ルート	投与時間																												
		ノバントロン	7 mg/m ²	点滴(側管)	30分	●	●	●																									
注射	4	商品名	用量	投与ルート	投与時間																												
		サンラビン	200 mg/m ²	点滴(側管)	3時間	●	●	●	●	●																							
		注射用水	10mg/mlで溶解																														
注射	5	商品名	用量	投与ルート	投与時間																												
		生理食塩液	50 ml	点滴(側管)	ルートフラッシュ	●	●	●	●	●																							
注射	6																																
注射	7																																
注射	8																																
注射	9																																
注射	10																																
内服	1																																
内服	2																																
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

70歳以上ではノバントロンを5mg/m²に減量する
前治療の影響がなくなり、十分に骨髄機能が回復した後に施行

減量規定

薬品名	ノバントロン	サンラビン			
標準投与量	7 mg/m ²	200 mg/m ²			
1段階減量	5 mg/m ²	150 mg/m ²			
2段階減量	3.5 mg/m ²	100 mg/m ²			
3段階減量					