

治療計画書

療法名 アザシチジン皮下注+ベネクレスタ(2サイクル目以降)

療法

1サイクル 4 週

疾患名 急性骨髄性白血病(AML)

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																												
			用量(/日)			用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
		照射部位	照射量																														
注射	1	グラニセトロン	3 mg	点滴	30分																												
		デキサート	6.6 mg			●	●	●	●	●	●																						
注射	2	ビダーザ	75 mg/m ²	皮下注																													
		注射用水	25mg/mlで溶解			●	●	●	●	●	●																						
注射	3																																
注射	4																																
注射	5																																
注射	6																																
注射	7																																
注射	8																																
注射	9																																
注射	10																																
内服	1	ベネクレスタ	400 mg	1日1回食後		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●				
内服	2																																
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

ベネクレスタ: 中程度以上のCYP3A阻害薬と併用する場合は減量を考慮。(詳細は添付文書参考)
適宜デカドロンの内服を追加する

減量規定

薬品名	ビダーザ				
標準投与量	75 mg/m ²				
1段階減量	37.5 mg/m ²				
2段階減量	25 mg/m ²				
3段階減量	中止				