

治療計画書

療法名 アザシチジン静注+ベネクレクスタ(1サイクル目)

療法

1サイクル 4 週

疾患名 急性骨髄性白血病(AML)

注射 内服	Rp	商品名	用量(回) 用量(日)	投与ルート 用法	投与時間																												
						照射部位 照射量																											
放射線		生理食塩液	100 ml	点滴	キープ	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
注射	1					●	●	●	●	●	●	●																					
注射	2	グラニセトロン デキサート	3 mg 6.6 mg	点滴(側管)	30分	●	●	●	●	●	●	●																					
注射	3	ビダーザ 注射用水 生理食塩液	75 mg/m ² 10mg/mlで溶解 50 ml	点滴(側管)	10分	●	●	●	●	●	●	●																					
注射	4	生理食塩液	50 ml	点滴(側管)	ルートフラッシュ	●	●	●	●	●	●	●																					
注射	5																																
注射	6																																
注射	7																																
注射	8																																
注射	9																																
注射	10																																
内服	1	ベネクレクスタ	100 mg	1日1回食後		●																											
内服	2	ベネクレクスタ	200 mg	1日1回食後			●																										
内服	3	ベネクレクスタ	400 mg	1日1回食後			●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

ベネクレクスタ: 中程度以上のCYP3A阻害薬と併用する場合は減量を考慮。(詳細は添付文書参考)
適宜デカドロンの内服を追加する

減量規定

薬品名	ビダーザ			
標準投与量	75 mg/m ²			
1段階減量	37.5 mg/m ²			
2段階減量	25 mg/m ²			
3段階減量	中止			