

## 治療計画書

療法名 JALSG ALL-97 地固め・強化(第1コース)

療法

1サイクル 日

疾患名 急性リンパ性白血病

注射 内服 放射線	Rp	商品名	用量(/回) 用量(/日)	投与ルート 用法	投与時間	1																													
							2																												
		照射部位	照射量			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
注射		生理食塩液	200 ml	点滴	キープ	●																													
注射	1																																		
注射		グラニセトン	3 mg	点滴(側管)	30分	●																													
注射	2																																		
注射		オンコピン 生理食塩液	※1.3 mg/m <sup>2</sup> 100 ml ※最大2mg	点滴(側管)	30分	●																													
注射	3																																		
注射		エンドキサン 生理食塩液	1000 mg/m <sup>2</sup> 500 ml	点滴(側管)	3時間	●																													
注射	4																																		
注射		ドキシルピシン 生理食塩液	60 mg/m <sup>2</sup> 500 ml	点滴(側管)	24時間	●																													
注射	5																																		
注射	6																																		
注射	7																																		
注射	8																																		
注射	9																																		
注射	10																																		
内服	1	プレドニゾン	60 mg/m <sup>2</sup>	1日1(-2)回朝(昼)食後		●●●																													
内服	2																																		
内服	3																																		
内服	4																																		
内服	5																																		
内服	6																																		
内服	7																																		
内服	8																																		
内服	9																																		
内服	10																																		
放射線			Gy																																

特記事項

JALSG ALL-97 髄注(DEXA+MTX)療法併用

減量規定

薬品名					
標準投与量					
1段階減量					
2段階減量					
3段階減量					