

治療計画書

療法名 JALSG ALL-97 地固め・強化(第2コース)

療法 1サイクル _____ 日

疾患名 急性リンパ性白血病

注射 内服	Rp	商品名	用量(/回) 用量(/日)	投与ルート 用法	投与時間	日																											
						1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
放射線		照射部位	照射量																														
注射	1	生理食塩液	200 ml	点滴	キープ		●	●																									
注射	2	グラニセトン	3 mg	点滴(側管)	30分		●	●																									
				①	9:30(day2)																												
					10:00(day3)																												
注射	3	メソトレキセート	50 mg/m ²	点滴(側管)	30分																												
		5%糖液	50 ml	全量50mlに調製	10:00		●																										
				①																													
注射	4	メソトレキセート	450 mg/m ²	点滴(側管)	23時間30分																												
		5%糖液	500 ml	全量500mlに調製	10:30		●																										
				①																													
注射	5	オンコビン	※1.3 mg/m ²	点滴(側管)	30分																												
		生理食塩液	100 ml	①	10:30			●																									
				※最大2mg																													
注射	6	ドキソルビシン	45 mg/m ²	点滴(側管)	30分																												
		生理食塩液	100 ml	①	11:00			●																									
注射	7	生理食塩液	50 ml	点滴(側管)	ルートフラッシュ																												
				①	11:30			●																									
注射	8	YDソリター-T3	500 ml	点滴	4時間			●	●	●	●																						
		7%メイロン	20 ml	②	2:00																												
注射	9	ロイコボリン	15 mg	静注(側管)					●	●																							
				②	4:00																												
注射	10	YDソリター-T3	500 ml	点滴	4時間		●	●	●	●																							
		7%メイロン	20 ml	②	6:00																												
注射	11	YDソリター-T3	500 ml	点滴	4時間		●	●	●	●																							
		7%メイロン	20 ml	②	10:00																												
注射	12	ダイアモックス	250 mg	静注(側管)				●	●	●	●																						
		生理食塩液	20 ml	②	10:00																												
注射	13	ロイコボリン	15 mg	静注(側管)					●	●																							
				②	10:00																												
注射	14	YDソリター-T3	500 ml	点滴	4時間		●	●	●	●																							
		7%メイロン	20 ml	②	14:00																												
注射	15	ロイコボリン	15 mg	静注(側管)					●	●																							
				②	16:00																												
注射	16	YDソリター-T3	500 ml	点滴	4時間		●	●	●																								
		7%メイロン	20 ml	②	18:00																												
注射	17	YDソリター-T3	500 ml	点滴	4時間		●	●	●	●																							
		7%メイロン	20 ml	②	22:00																												
注射	18	ダイアモックス	250 mg	静注(側管)			●	●	●	●																							
		生理食塩液	20 ml	②	22:00																												
注射	19	ロイコボリン	15 mg	静注(側管)					●	●																							
				②	22:00																												
内服	1	プレドニゾロン	60 mg/m ²	1日1(-2)回朝(昼)食後			●	●	●																								

特記事項

- ・JALSG ALL-97 髄注(DEXA+MTX)療法併用
- ・ロイコボリンはメソトレキセート投与開始36時間後より6時間毎に8回投与

減量規定

薬品名	標準投与量	1段階減量	2段階減量	3段階減量