

治療計画書

療法名

JALSG ALL-202地固め2(C2)

療法

1サイクル _____ 日

疾患名 Ph染色体陽性急性リンパ性白血病(15~64歳)

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
						用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
注射	1	メソレキセート	15 mg	髄注																													
		キロサイド	40 mg			●																											
		デキサート	3.3 mg																														
		生理食塩液	5 ml	全量5mlに調製																													
注射	2																																
注射	3																																
注射	4																																
注射	5																																
注射	6																																
注射	7																																
注射	8																																
注射	9																																
注射	10																																
内服	1	グリベック	600 mg	1日1回食後		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●			
内服	2																																
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

- 寛解後療法
- C1(地固め療法1)とC2を1サイクルとして4サイクル施行

減量規定

薬品名	標準投与量	1段階減量	2段階減量	3段階減量