

## 治療計画書

療法名

アザシチジン静注

療法

1サイクル 4 週

疾患名 骨髄異形成症候群

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																																			
			用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28								
		照射部位	照射量																																					
注射	1	生理食塩液	100 ml	点滴	キープ	●	●	●	●	●																														
注射	2	グラニセトロン デキサート	3 mg 6.6 mg	点滴(側管)	30分	●	●	●	●	●																														
注射	3	ビダーザ 注射用水 生理食塩液	75 mg/m <sup>2</sup> 10mg/mlで溶解 50 ml	点滴(側管)	10分	●	●	●	●	●																														
注射	4	生理食塩液	50 ml	点滴(側管)	ルートフラッシュ	●	●	●	●	●																														
注射	5																																							
注射	6																																							
注射	7																																							
注射	8																																							
注射	9																																							
注射	10																																							
内服	1																																							
内服	2																																							
内服	3																																							
内服	4																																							
内服	5																																							
内服	6																																							
内服	7																																							
内服	8																																							
内服	9																																							
内服	10																																							
放射線			Gy																																					

## 特記事項

- ・月～金曜日に投与(休日をはさみ4日と3日に分かれる場合あり)
- ・末梢ルートの確保が困難な症例では、皮下投与へ変更可能
- ・適宜デカドロンの内服を追加する

## 減量規定

薬品名	ビダーザ				
標準投与量	75 mg/m <sup>2</sup>				
1段階減量	37.5 mg/m <sup>2</sup>				
2段階減量	25 mg/m <sup>2</sup>				
3段階減量	中止				