

治療計画書

療法名

R-VeMP

療法

1サイクル

4週

疾患名 再発非ホジキンリンパ腫(6サイクル)

| 注射 内服 放射線 | Rp | 商品名 | 用量(/回) 用量(/日) | 投与ルート 用法 | 投与時間 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|-----------------|----|-----------------|---------------------------------|------------------------|-----------|---|---|---|---|---|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
| | | | | | | 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | 照射部位 | 照射量 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 1 | 生理食塩液 | 200 ml | 点滴 | キープ | ● | ● | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 2 | リツキシマブ 生理食塩液 | 375 mg/m ² ml | 点滴(側管) 10倍希釈 | フローシードに従う | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 3 | プレドニン 生理食塩液 | 60 mg/m ² 100 ml | 点滴(側管) (day4、5はメイン) | 30分 | ● | ● | ● | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 4 | グラニセトン | 3 mg | 点滴(側管) | 30分 | ● | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 5 | ノバントロン 生理食塩液 | 6 mg/m ² 100 ml | 点滴(側管) | 30分 | ● | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 6 | ラスデット 生理食塩液 | 100 mg/m ² 500 ml | 点滴(側管) | 2時間 | ● | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 7 | 生理食塩液 | 50 ml | 点滴(側管) | ルートフラッシュ | ● | ● | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 1 | ネオマレルミンTR | 6 mg | リツキシマブ投与30分前 | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 2 | ブルフェン | 200 mg | リツキシマブ投与30分前 | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 放射線 | | | Gy | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

特記事項

・リツキシマブの最大投与回数8回

減量規定

| 薬品名 | 標準投与量 | 1段階減量 | 2段階減量 | 3段階減量 |
|-----|-------|-------|-------|-------|
| | | | | |
| | | | | |
| | | | | |