

治療計画書

療法名

GPD

(ID 0) ☆7

療法 1サイクル 4 週

疾患名 再発・難治性非ホジキンリンパ腫(3~6サイクル)

注射 内服 放射線	Rp	商品名	用量(/回) 用量(/日)	投与ルート 用法	投与時間																														
						照射部位 照射量																													
						1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
注射	1	生理食塩液	200 ml	点滴	キープ																														
注射	2	ソリューゲンF	500 ml	点滴(側管)	2時間																														
注射	3	グラニセトロン	3 mg	点滴(側管)	30分																														
注射	4	デキサート 生理食塩液	6.6 mg 100 ml	点滴(側管)	30分																														
注射	5	ゲムシタピン 生理食塩液	1000 mg/m ² 100 ml	点滴(側管)	30分																														
注射	6	生理食塩液	500 ml	点滴(側管)	2時間																														
注射	7	シスプラチン 生理食塩液	75 mg/m ² 500 ml	点滴(側管) 全量500mlに調製	2時間																														
注射	8	ソリューゲンF	500 ml	点滴(側管)	2時間																														
注射	9	ソリューゲンF	500 ml	点滴(側管)	2時間																														
注射	10	生理食塩液	50 ml	点滴(側管)	ルートフラッシュ																														
内服	1	デカドロン	40 mg	1日1(-2)回朝(昼)食後		●	●	●	●																										
内服	2	イメンド	125 mg	抗癌剤の1時間-1時間半前		●																													
内服	3	イメンド	80 mg	1日1回朝食後		●	●																												
内服	4																																		
内服	5																																		
内服	6																																		
内服	7																																		
内服	8																																		
内服	9																																		
内服	10																																		
放射線			Gy																																

特記事項

ラシックス注は適宜使用

減量規定

薬品名	ゲムシタピン	シスプラチン		
標準投与量	1000 mg/m ²	75 mg/m ²		
1段階減量	500 mg/m ²	37.5 mg/m ²		
2段階減量	中止	中止		
3段階減量				