

治療計画書

療法名

アドセトリス

療法

1サイクル 3 週間

疾患名 再発又は難治性のCD30陽性の末梢性T細胞リンパ腫

注射 内服 放射線	Rp	商品名	用量(回)	投与ルート	投与時間																																
			用量(日)	用法		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28				
放射線		照射部位	照射量																																		
注射	1	生理食塩液	200 ml	点滴	キープ	●																															
注射	2	デキサート 生理食塩液	6.6 mg 100 ml	点滴(側管)	30分	●																															
注射	3	生理食塩液	50 ml	点滴(側管)	10分	●																															
注射	4	アドセトリス 注射用水 生理食塩液	1.8 mg/kg 5mg/mlで溶解 ※0.4-1.2mg/mlで希釈 ※最大180mg	点滴(側管)	30分	●																															
注射	5	生理食塩液	50 ml	点滴(側管)	15分	●																															
注射	6																																				
注射	7																																				
注射	8																																				
注射	9																																				
注射	10																																				
内服	1																																				
内服	2																																				
内服	3																																				
内服	4																																				
内服	5																																				
内服	6																																				
内服	7																																				
内服	8																																				
内服	9																																				
内服	10																																				
放射線			Gy																																		

特記事項

※アドセトリスが50mg以下の場合は生理食塩液50mlで、50-150mgの場合は生理食塩液100mlで、150mg以上の場合は250mlで希釈する。
CCr30未満で減量を考慮

減量規定

薬品名	アドセトリス			
標準投与量	1.8 mg/kg			
1段階減量	1.2 mg/kg			
2段階減量	中止			
3段階減量				