

治療計画書

療法名

G-CVP(1サイクル目)

療法

1サイクル 3 週

(ID 0) ☆7

疾患名 CD20陽性の濾胞性リンパ腫(8サイクル)

注射 内服 放射線	Rp	商品名	用量(/回) 用量(/日)	投与ルート 用法	投与時間																											
						照射部位	照射量	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
注射	1	生理食塩液	200 ml	点滴	キープ																											
注射	2	プレドニン	100 mg	点滴(側管)	30分																											
		生理食塩液	50 ml																													
注射	3	生理食塩液	50 ml	点滴(側管)	30分																											
注射	4	ガザイバ	1000 mg/body	点滴(側管)	フローシートに従う																											
		生理食塩液	210 ml	全量250mlに調製																												
注射	5	生理食塩液	50 ml	点滴(側管)	ルートフラッシュ																											
注射	6	グラニセトロン	3 mg	点滴(側管)	30分																											
注射	7	オンコピン	※1.4 mg/m ²	点滴(側管)	30分																											
		生理食塩液	100 ml																													
注射	8	エンドキサン	750 mg/m ²	点滴(側管)	2時間																											
		生理食塩液	500 ml																													
注射	9																															
注射	10																															
内服	1	ネオマレルミンTR	6 mg	ガザイバ投与30分前																												
内服	2	カロナール	1000 mg	ガザイバ投与30分前																												
内服	3	プレドニゾン	100 mg	1日1(-2)回朝(昼)食後																												
内服	4																															
内服	5																															
内服	6																															
内服	7																															
内服	8																															
内服	9																															
内服	10																															
放射線			Gy																													

特記事項

ガザイバ:0. 2又は0. 22μ mのインラインフィルターを使用すること。

減量規定

薬品名	標準投与量	1段階減量	2段階減量	3段階減量