

治療計画書

療法名

C-MOPP+ABVD(C-MOPPコース)

療法

1サイクル

4 週

疾患名 ホジキンリンパ腫(※5サイクル)

注射 内服	Rp 放射線	商品名	用量(回) 用量(日)	投与ルート 用法	投与時間	日																											
						1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
		照射部位	照射量																														
1	注射	生理食塩液	200 ml	点滴	キープ	●											●																
2	注射	グラニセトロン	3 mg	点滴(側管)	30分	●											●																
3	注射	オンコピン	※1.4 mg/m ²	点滴(側管)	30分	●											●																
		生理食塩液	100 ml			●											●																
			※最大2mg																														
4	注射	エンドキサソ	550 mg/m ²	点滴(側管)	2時間	●											●																
		生理食塩液	500 ml			●											●																
5	注射																																
6	注射																																
7	注射																																
8	注射																																
9	注射																																
10	注射																																
1	内服	プロカルバジン	100 mg/m ²	1日1回朝食後		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●												
2	内服	プレドニゾン	40 mg/m ²	1日1(-2)回朝(昼)食後		●	●	●									●	●	●														
3	内服																																
4	内服																																
5	内服																																
6	内服																																
7	内服																																
8	内服																																
9	内服																																
10	内服																																
放射線			Gy																														

特記事項

※C-MOPPとABVDを交互に5回繰り返す
・プロカルバジンに対する制吐療法は適宜選択する

減量規定

薬品名					
標準投与量					
1段階減量					
2段階減量					
3段階減量					