

治療計画書

療法名

Kd

療法

1サイクル 4 週

疾患名 再発又は難治性の多発性骨髄腫(1サイクル目)

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																																			
			用量(/日)	用法		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28							
放射線		照射部位	照射量																																					
注射	1	5%糖液	250 ml	点滴	キープ	●	●																																	
注射	2	5%糖液	50 ml	点滴(側管)	キープ	●	●																																	
注射	3	カイプロリス 注射用水 5%糖液	※20 mg/m ² 2mg/mlで溶解 100 ml	点滴(側管)	30分間	●	●																																	
注射	4	カイプロリス 注射用水 5%糖液	※56 mg/m ² 2mg/mlで溶解 100 ml	点滴(側管)	30分間																																			
注射	5	5%糖液	50 ml	点滴(側管)	ルートフラッシュ	●	●																																	
注射	6																																							
注射	7																																							
注射	8																																							
注射	9																																							
注射	10																																							
内服	1	レナデックス	20 mg	1日1(-2)回朝(昼)食後		●	●																																	
内服	2																																							
内服	3																																							
内服	4																																							
内服	5																																							
内服	6																																							
内服	7																																							
放射線			Gy																																					

特記事項

- ・血液毒性またはGrade3以上の非血液毒性が出現した場合には休薬し、症状が回復した場合は以下を目安としてカイプロリスの減量を考慮する。
- ・CCr<15ml/minとなった場合はカイプロリスを休薬し、CCr≥15以上に回復した場合は再開を検討する。

減量規定

薬品名	カイプロリス	レナデックス		
標準投与量	56 mg/m ²	20 mg		
1段階減量	45 mg/m ²	12 mg		
2段階減量	36 mg/m ²	8 mg		
3段階減量	27 mg/m ²	中止		
4段階減量	中止			