

疾患名 手術不能又は再発大腸がん

| 注射 内服 放射線 | Rp | 商品名 | 用量(/回) 用量(/日) | 投与ルート 用法 | 投与時間 | 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|-----------------|----|-----------------|-----------------------|----------------|------|--|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|--|--|--|--|
| | | | | | | 照射部位 照射量 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 1 | グラニセトロン | 3 mg | 点滴 | 30分 | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | デキサート | 6.6 mg | | | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 2 | トボテシン | 150 mg/m ² | 点滴 | 2時間 | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | 生理食塩液 | 500 ml | | | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 3 | 生理食塩液 | 50 ml | 点滴 | 15分 | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 注射 | 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 1 | 半夏瀉心湯 | 7.5 g | 1日3回食間/食前 | | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | | | | |
| 内服 | 2 | ウルソ | 300 mg | 1日3回食後 | | ● | ● | ● | | | | | | | | | | | | | | | | ● | ● | ● | | | | | | | |
| 内服 | 3 | プリンペラン | 15 mg | 1日3回食前 | | ● | ● | ● | | | | | | | | | | | | | | | | ● | ● | ● | | | | | | | |
| 内服 | 4 | 重曹 | 3 g | 1日3回食間 | | ● | ● | ● | | | | | | | | | | | | | | | | ● | ● | ● | | | | | | | |
| 内服 | 5 | 酸化マグネシウム/マグラックス | 1.2-1.8 g | 1日3回食後 | | ● | ● | ● | | | | | | | | | | | | | | | | ● | ● | ● | | | | | | | |
| 内服 | 6 | デカドロン | 8 mg | 1日1(-2)回朝(昼)食後 | | ● | ● | | | | | | | | | | | | | | | | | ● | ● | | | | | | | | |
| 内服 | 7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 内服 | 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 放射線 | | | Gy | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

特記事項

※UGT1A1遺伝子多型にてホモ接合体、複合ヘテロ接合体をもつ場合は、1段階以上の減量を行う(最大50%まで減量を考慮)。

減量規定

| 薬品名 | 標準投与量 | 1段階減量 | 2段階減量 | 3段階減量 |
|-----|-------|-------|-------|-------|
| | | | | |
| | | | | |
| | | | | |