

疾患名 手術不能又は再発の直腸・結腸がん

注射	商品名	用量(/回)	投与ルート	投与時間	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
		内服	Rp	用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
放射線	照射部位		照射量																													
注射	1	グラニセトロン	3 mg	点滴	30分																											
注射	1	デキサート	6.6 mg			●															●											
注射	2	トポテシン	125 mg/m ²	点滴	2時間																											
注射	2	生理食塩液	500 ml			●															●											
注射	3	生理食塩液	50 ml	点滴	15分																	●										
注射	4																															
注射	5																															
注射	6																															
注射	7																															
注射	8																															
注射	9																															
注射	10																															
内服	1	TS-1	80 mg/m ²	1日2回朝夕食後		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●											
内服	2	半夏瀉心湯	7.5 g	1日3回食間/食前		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
内服	3	ウルソ	300 mg	1日3回食後		●	●	●													●	●	●									
内服	4	プリンペラン	15 mg	1日3回食前		●	●	●												●	●	●										
内服	5	重曹	3 g	1日3回食間		●	●	●											●	●	●											
内服	6	酸化マグネシウム/マグラックス	1.2-1.8 g	1日3回食後		●	●	●												●	●	●										
内服	7	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●													●	●	●										
内服	8																															
内服	9																															
内服	10																															
放射線			Gy																													

特記事項

・76歳以上:1段階減量より開始する。

・セカンドラインとして使用

※UGT1A1遺伝子多型にてホモ接合体、複合ヘテロ接合体をもつ場合は、1段階以上の減量を行う(最大50%まで減量を考慮)。

減量規定

薬品名	トポテシン※	TS-1(1.25m ² 未満)	TS-1(1.25m ² 以上1.5m ² 未満)	TS-1(1.5m ² 以上)	
標準投与量	125 mg/m ²	80 mg/day	100 mg/day	120 mg/day	
1段階減量	100 mg/m ²		80 mg/day	100 mg/day	
2段階減量	80 mg/m ²			80 mg/day	
3段階減量					