

治療計画書

療法名

SOX

療法

1サイクル 3 週

疾患名 治療切除不能な進行・再発の結腸・直腸がん

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																												
			用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
		照射部位	照射量																														
注射	1	グラニセトロン	3 mg	点滴	30分																												
		デキサート	6.6 mg			●																											
注射	2	エルブラット	130 mg/m ²	点滴	2時間																												
		5%糖液	250 ml			●																											
注射	3	生理食塩液	50 ml	点滴	15分																												
注射	4																																
注射	5																																
注射	6																																
注射	7																																
注射	8																																
注射	9																																
注射	10																																
内服	1	TS-1※	80 mg/m ²	1日2回朝夕食後		●●●●●●●●●●●●●●●●●●●●●●●●●●																											
内服	2	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●●																											
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

エルブラット総投与量が600mg/m²を超えた場合、エルブラットは担当医の判断で適宜休止して次のサイクルを開始する。
 制吐剤としてデカドロン錠を適宜追加する(day2~8mgを2(~3)日)。
 ※1日目の夕食後より服用開始。

減量規定

薬品名	エルブラット	TS-1(1.25m ² 未満)	TS-1(1.25m ² 以上1.50m ² 未満)	TS-1(1.50m ² 以上)	
標準投与量	130 mg/m ²	80 mg/day	100 mg/day	120 mg/day	
1段階減量	100 mg/m ²	60 mg/day	80 mg/day	100 mg/day	
2段階減量	75 mg/m ²	50 mg/day	60 mg/day	80 mg/day	
3段階減量					