

疾患名 治癒切除不能な進行・再発の結腸・直腸がん

注射	Rp	商品名	用量(回)	投与ルート	投与時間																															
						用量(日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
放射線		照射部位	照射量																																	
注射	1	グラニセトロン デキサート	3 mg 6.6 mg	点滴	30分	●																														
注射	2	生理食塩液	50 ml	点滴	15分	●																														
注射	3	アバステン 生理食塩液	7.5 mg/kg 100 ml	点滴	※初回90分 2回目60分可 3回目以降 30分可	●																														
注射	4	生理食塩液	50 ml	点滴	15分	●																														
注射	5	エルプラット 5%糖液	130 mg/m ² 250 ml	点滴	2時間	●																														
注射	6	生理食塩液	50 ml	点滴	15分	●																														
注射	7																																			
注射	8																																			
注射	9																																			
注射	10																																			
内服	1	TS-1※※	80 mg/m ²	1日2回朝夕食後		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
内服	2	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●																													
内服	3																																			
内服	4																																			
内服	5																																			
内服	6																																			
内服	7																																			
内服	8																																			
内服	9																																			
内服	10																																			

特記事項

エルプラット総投与量が600mg/m²を超えた場合、エルプラットは担当医の判断で適宜休止して次のサイクルを開始する。

制吐剤としてデカドロン錠を適宜追加する(day2~8mgを2(~3)日)。

※アバステンは初回90分で投与、忍容性が良ければ2回目以降60分、3回目以降30分で投与可。

※※1日目の夕食後より服用開始。

減量規定

薬品名	アバステン	エルプラット	TS-1(1.25m ² 未満)	TS-1(1.25m ² 以上1.50m ² 未満)	TS-1(1.50m ² 以上)
標準投与量	7.5 mg/kg	130 mg/m ²	80 mg/day	100 mg/day	120 mg/day
1段階減量	減量なし	100 mg/m ²	60 mg/day	80 mg/day	100 mg/day
2段階減量		75 mg/m ²	50 mg/day	60 mg/day	80 mg/day
3段階減量					