

治療計画書

療法名

TAS-102+アバスチン

療法

1サイクル

4 週

疾患名 治癒切除不能な進行・再発の結腸・直腸がん

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間																													
			用量(/日)	用法		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
		照射部位	照射量																															
注射	1	生理食塩液	50 ml	点滴	15分																													
						●																												
注射	2	アバスチン	5 mg/kg	点滴	※初回90分																													
		生理食塩液	100 ml		2回目60分可																													
					3回目以降	●																												
注射	3	生理食塩液	50 ml	点滴	15分																													
						●																												
注射	4																																	
注射	5																																	
注射	6																																	
注射	7																																	
注射	8																																	
注射	9																																	
注射	10																																	
内服	1	TAS-102	下記参照	1日2回朝夕食後		●	●	●	●	●	●	●	●	●	●	●																		
内服	2																																	

特記事項

TAS-102の減量は、下記表の初回投与量から10mg/日ずつ減量し、最低投与量は、40mg/m²/日。

TAS-102 減量基準	1日投与量
	初回投与量
1.07m ² 未満	70mg/日
1.07m ² 以上1.23m ² 未満	80mg/日
1.23m ² 以上1.38m ² 未満	90mg/日
1.38m ² 以上1.53m ² 未満	100mg/日
1.53m ² 以上1.69m ² 未満	110mg/日
1.69m ² 以上1.84m ² 未満	120mg/日
1.84m ² 以上1.99m ² 未満	130mg/日
1.99m ² 以上2.15m ² 未満	140mg/日
2.15m ² 以上	150mg/日

減量規定

薬品名				
標準投与量				
1段階減量				
2段階減量				
3段階減量				