

治療計画書

療法名

ゲムシタビン+シスプラチン

療法

1サイクル 3 週

疾患名 胆道がん(8サイクル)

注 射	Rp	商品名	用量(/回)	投与ルート	投与時間	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
						用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
放射線		照射部位	照射量																														
注 射	1	ソリューゲンF	500 ml	点滴	2時間																												
						●																											
注 射	2	ソリューゲンF	500 ml	点滴	2時間																												
						●																											
注 射	3	グラニセトロン	3 mg	点滴	30分																												
		デキサート	9.9 mg																														
注 射	4	ゲムシタビン	1000 mg/m ²	点滴	30分																												
		生理食塩液	100 ml																														
注 射	5	生理食塩液	50 ml	点滴	15分																												
						●																											
注 射	6	シスプラチン	25 mg/m ²	点滴	1時間																												
		生理食塩液	250 ml	全量250mlに調製																													
注 射	7	ソリューゲンF	500 ml	点滴	2時間																												
						●																											
注 射	8	ソリューゲンF	500 ml	点滴	2時間																												
						●																											
注 射	9																																
注 射	10																																
内服	1	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●	●																									
内服	2																																
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

減量規定

薬品名	ゲムシタビン	シスプラチン	
標準投与量	1000 mg/m ²	CCr>60	100 %
1段階減量	800 mg/m ²	45<CCr≤60	75 %
2段階減量	600 mg/m ²	CCr≤45	中止
3段階減量			