

治療計画書

療法名

ゲムシタビン+タルセバ

(ID 0) ☆7

療法 1サイクル 4 週

疾患名 治癒切除不能な腭癌

注射 内服 放射線	Rp	商品名	用量(/回) 用量(/日)	投与ルート 用法	投与時間	1																						
						1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
注射	1	グラニセトロン	3 mg	点滴	30分																							
		デキサート	6.6 mg				●																					
注射	2	ゲムシタビン	1000 mg/m ²	点滴	30分																							
		生理食塩液	100 ml				●																					
注射	3	生理食塩液	50 ml	点滴	15分																							
							●																					
注射	4																											
注射	5																											
注射	6																											
注射	7																											
注射	8																											
注射	9																											
注射	10																											
内服	1	タルセバ	100 mg	1日1回空腹時		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
内服	2																											
内服	3																											
内服	4																											
内服	5																											
内服	6																											
内服	7																											
内服	8																											
内服	9																											
内服	10																											
放射線			Gy																									

特記事項

グラニセトロンは適宜使用

減量規定

薬品名	ゲムシタビン	タルセバ			
標準投与量	1000 mg/m ²	100 mg			
1段階減量	750 mg/m ²	50 mg			
2段階減量					
3段階減量					