

# 治療計画書

療法名

TC(ドセタキセル+シクロホスファミド)

療法

1サイクル

3 週

進行・再発乳がん

疾患名  乳がんの術後補助療法(4サイクル)

注射 内服 放射線	Rp	商品名	用量(/回) 用量(/日)	投与ルート 用法	投与時間																												
						1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
照射部位		照射量																															
注射	1	生理食塩液	20 ml	CVCポート	ルートキーブ	●																											
注射	2	グラニセトロン	3 mg	点滴	30分	●																											
		ファモチジン	20 mg																														
		デキサート	6.6 mg																														
注射	3	ドセタキセル	75 mg/m <sup>2</sup>	点滴(側管)	1時間	●																											
		生理食塩液	250 ml																														
注射	4	生理食塩液	50 ml	点滴	15分	●																											
注射	5	エンドキサン	600 mg/m <sup>2</sup>	点滴(側管)	1時間	●																											
		生理食塩液	100 ml																														
注射	6	生理食塩液	50 ml	点滴	15分	●																											
注射	7	ヘパリンNaロックシリンジ	100 IU	CVCポート		●																											
注射	8																																
注射	9																																
注射	10																																
内服	1	レスタミン	50 mg	ドセタキセルの30分前まで		●																											
内服	2	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●●																											
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

減量規定

薬品名	ドセタキセル	エンドキサン			
標準投与量	75 mg/m <sup>2</sup>	600 mg/m <sup>2</sup>			
1段階減量	60 mg/m <sup>2</sup>	500 mg/m <sup>2</sup>			
2段階減量	45 mg/m <sup>2</sup>	400 mg/m <sup>2</sup>			
3段階減量	中止	中止			