治療計画書 療法名 カルボプラチン(3週毎)+パクリタキセル+キイトルーダ(3週毎) 療法 1サイクル 3 週

_{疾患名} ホルモン受容体陰性かつHER2陰性で再発高リスクの乳癌における術前薬物療法(4コース)

注射			用量(/回)	投与ルート	投与時間	1						2					3						4			Т	П
内服	Rn	商品名	用量(/日)	用法		1	2	3 4	4 5	6	7		10	11	2 13			6 1	7 18	19	20			24	25 2	6 27	28
放射線		照射部位	照射量	71372	•	Ė	-			-	ť		10	÷	2 10	-	10		, 10	10	20		- 20	-	20 2	-	-
从对外		生理食塩液		CVCポート																			+	Н	-	-	H
		<u> </u>	20 ml	CVC/V-L	ルートキーノ	4																		Н			
注射	1					•						•												Н			
射																								Н			
																								Ш			
		キイトルーダ	200 mg/body	点滴	30分																						П
注		生理食塩液	100 ml																								
注射	2					•																		Н			
733						1																		Ш			
		生理食塩液	50 ml	点滴	15分				+	+		+	+			_	+	+	+	H		+	+	H	-	+	Н
:+		工生及塩水	30 1111	杰旭	1371	1																		Ш			
注射	3					•																		Н			
别						4																		Н			
		/×=	_		11																		_	Н			Ш
		グラニセトロン	3 mg	点滴	30分																			Н			
注		ファモチジン	20 mg			•																		Н			
射		デキサート	9.9 mg			•																		Ш			
		アロカリス	235 mg																					Ш			
		生理食塩液	50 ml	点滴	15分																			П			П
注	Ιİ					1_																		Н			
注射	5					•																					
21																											
	H	ファモチジン	20 mg	点滴	30分		H		+	۰	\blacksquare	+	H		+	+	-	+	+	H	\dashv	+	+	\forall	+	+	\forall
24					3077	-																		Н			
注射	6	デキサート	9.9 mg																					Н			
射		生理食塩液	50 ml			4																		Н			
																								Ш			
		パクリタキセル	80 mg/ m i	点滴	1時間																			Н			
注射	,	生理食塩液	250 ml																					Н			
射						•																		Н			
																								Н			
		生理食塩液	50 ml	点滴	15分							1					1						\top	Ħ	_	+	П
注				211171-3	,,	1																		Н			
注射	8																							Н			
划						-																		Н			
		L :: _1^	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	F 144	4 0+ 88																		+	Н	_	+	Н
		カルボプラチン	XAUC5	点滴	1時間	4																		Н			
注射	9	生理食塩液	250 ml			•																		Н			
射			※最大750mg																					Н			
																								Н			
		生理食塩液	50 ml	点滴	15分																			П			П
注						1_																		Н			
注射	10					•																		Н			
~,,						1																		Н			
		へパリンNaロックシリンジ	100 IU	CVCポート												-				Н			+	H	+	+	Н
:+		· · · · · · · · · · · · · · · · · · ·	100 10	0,00/1		-																		Н			
注射	10					•						•												Н			
别						4																		Н			
	Ц			0.6-11.5			Щ			_			Ш			_					Щ	\downarrow	\perp	\sqcup			Щ
		レスタミン	50 mg	パクリタキセルの		lacktriangle						•					•							Ш			Ш
内服	2	デカドロン	4 mg	1日1回朝	食後		lacktriangle	•	•)															Ш			
内服	3																							Ш			
内服	4																										П
内服	5												П											П			П
内服	-																						T	П			П
放射線			Gy													_				Н			+	Н			П
	ш		,		特記事項					_						!											\dashv
イン	51	'ンフィルター(0.22 μ m以下)を使	目すること		17107-7																						\dashv
	ر. ا	2 2 170 % (0.22 µ III M 1') & B	111 7 OCC0																								
					減量規定																						\dashv
遨	品	名 パクリタキセル	カルボプ	ラチン	<i>*</i> /ハ <u>エ</u> /ハレ						I								T								\dashv
本	护	<u> </u>	AUC								1								+								\dashv
		ブ重 30 mg/m 咸量 70 mg/ m ²	中山								1								+								\dashv
2段	<u>階</u> :	咸量 70 mg/ m 咸量 中止	7.1	_							1								+								\dashv
3段	陛:	ᇫᇫᅟᅟᅟᅮᄯ 咸量	+	+							+								+								\dashv
JAX	r# //	w±	<u> </u>								1								1_								