

疾患名 神経内分泌腫瘍

注射 内服 放射線	Rp	商品名	用量(/回)	投与ルート	投与時間	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
			用量(/日)			用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
		照射部位	照射量																														
注射	1	アロキシ	0.75 mg	点滴	30分																												
		デキサート	9.9 mg			●																											
		生理食塩液	100 ml																														
注射	2	デキサート	6.6 mg	点滴	30分																												
		生理食塩液	100 ml			●	●																										
注射	3	生理食塩液	500 ml	点滴	2時間																												
		硫酸Mg補正液	8 mEq			●																											
注射	4	生理食塩液	50 ml	点滴	15分																												
						●	●																										
注射	5	ラステット	100 mg/m ²	点滴	2時間																												
		5%光糖液	500 ml			●	●	●																									
注射	6	生理食塩液	500 ml	点滴	2時間																												
						●																											
注射	7	シスプラチン	80 mg/m ²	点滴	2時間																												
		生理食塩液	500 ml	全量500mlに調製		●																											
注射	8	YDソリター-T3	500 ml	点滴	2時間																												
		フロセミド	20 mg			●																											
注射	9	YDソリター-T3	500 ml	点滴	2時間																												
						●	●																										
注射	10	ソリューゲンF	500 ml	点滴	2時間																												
						●	●	●																									
内服	1	アプレピタント	125 mg	抗癌剤の1時間-1時間半前		●																											
内服	2	アプレピタント	80 mg	1日1回朝食後		●	●																										
内服	3	デカドロン	8 mg	1日1(-2)回朝(昼)食後				●	●																								
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
放射線			Gy																														

特記事項

減量規定

薬品名	エトポシド	シスプラチン		
標準投与量	100 mg/m ²	80 mg/m ²		
1段階減量	80 mg/m ²	60 mg/m ²		
2段階減量	60 mg/m ²	40 mg/m ²		