

# 治療計画書

療法名

**キイトルーダ(3週毎)+SOX**

療法

1サイクル **3** 週

疾患名 **治癒切除不能な進行・再発の胃癌**

注射 内服 放射線	Rp	商品名	用量(/回) 用量(/日)	投与ルート	投与時間	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28																											
						用法																											
放射線		照射部位	照射量																														
注射	1	生理食塩液	50 ml	点滴	15分																												
						●																											
注射	2	キイトルーダ	200 mg/body	点滴	30分																												
		生理食塩液	100 ml			●																											
注射	3	生理食塩液	50 ml	点滴	15分																												
						●																											
注射	4	グラニセトロン	3 mg	点滴	30分																												
		デキサート	6.6 mg			●																											
注射	5	エルプラット	130 mg/m <sup>2</sup>	点滴	2時間																												
		5%糖液	250 ml			●																											
注射	6	生理食塩液	50 ml	点滴	15分																												
						●																											
注射	7																																
注射	8																																
注射	9																																
注射	10																																
内服	1	S-1※	80 mg/m <sup>2</sup>	1日2回朝夕食後		●●●●●●●●●●●●●●●●●●●●●●●●●●●●																											
内服	2	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●●																											
内服	3																																
内服	4																																
内服	5																																
内服	6																																
内服	7																																
内服	8																																
内服	9																																
内服	10																																
放射線			Gy																														

特記事項

キイトルーダは最大35サイクルまで。  
 インラインフィルター(0.2~5µm)を使用すること。  
 ※S-1は1日目の夕食後より服用開始。

減量規定

薬品名	エルプラット	S-1(1.25m <sup>2</sup> 未満)	S-1(1.25m <sup>2</sup> 以上1.50m <sup>2</sup> 未満)	S-1(1.50m <sup>2</sup> 以上)
標準投与量	130 mg/m <sup>2</sup>	80 mg/day	100 mg/day	120 mg/day
1段階減量	100 mg/m <sup>2</sup>		80 mg/day	100 mg/day
2段階減量	75 mg/m <sup>2</sup>			80 mg/day
3段階減量	50 mg/m <sup>2</sup>			