治療計画書

療法名

キイトルーダ(3週毎)+XELOX

療法 1サイクル 3 週

疾患名 治癒切除不能な進行・再発の胃癌

34 64		Т	마루//리)	+n ⊢ ı	+ル ヒ 1±88	1 , 1	_		1				_		_	_		٦	_		_	_			_	_		$\overline{}$			
注射	_	商品名	用量(/回)	投与ルート 用法	技 子 時 间	1		3 4	_	H		2			4 .			3	16			0 0	0.	4	0.0	0.1					
内服 放射線			用量(/日)	用法		1	2	3 4	5	6	7	8	9 1	0 1	1 13	2 13	14	15	16	17 1	8 1	9 20	21	22	23	24 2	25 2	3 27	28		
放射線		照射部位	照射量	F 25	157		4			Н		4		_	_	-	Н		4	4	_	+		4		4		+-	▙		
:+		生理食塩液	50 ml	点滴	15分																										
注射	1																														
别																															
		キイトルーダ	200 mg/body	点滴	30分		+			Н		+	+	+	+		Н		\dashv		+			+				+	╆		
注射		キイトルーダ 生理食塩液	100 ml	W(VID)	0073	•																									
	2		100 1111																												
711																															
		生理食塩液	50 ml	点滴	15分														T		T			T				\top			
注射	ı																														
射	3					•																									
注射		グラニセトロン	3 mg	点滴	30分																										
	4	デキサート	6.6 mg																												
	4					•																									
注射		エルプラット	130 mg/m ²	点滴	2時間																										
	_	5%糖液	250 ml																												
	5																														
		生理食塩液	50 ml	点滴	15分												Н		\dashv					_				+	Н		
注	1			71111-3																											
注射	6																														
711																															
							+			Н		\dashv		+			H		\dashv	+	\dagger			7				+	H		
注						-																									
注射	7																														
711																															
							\dashv			H		+	+	+		+	H		\dashv		+	+		+		+		+			
注																															
注射	8					-	П																								
20						-																									
内服	1	カペシタビン※	2000 mg/m ²	1日2回朝会	│ ▽			• •											+	+	+	+		\dashv		+		+	Н		
中服	1	デカドロン	8 mg	1日1(-2)回朝			•												\dashv	+	+			\dashv	+	+		+	\vdash		
内服			o mg		(生/民)	H	_			Н	${oldsymbol{ert}}$	+		+			H		\dashv	+	+	-		\dashv		+	+	+	\vdash		
内服							_			Н		-					Н		\dashv		+			\dashv				+	⊢		
内服							4			Н	\vdash	+		+	+	H	Н		\dashv	+	+	+		\dashv		+	+	+	\vdash		
							4			Н	\vdash	+		+	+	H	Н		\dashv	+	+	+		\dashv		+	+	+	\vdash		
内服							4			Н	\square	4		+			Н		\dashv	-	+			4		-		_	₽		
内服							_			Н	\sqcup						Н		_					_				+			
内服									L	Ш		4		-			Ш		4		4			4		4		\perp	_		
内服	_									Ш							Ш		_									4	Щ		
内服	10				11 == =																								Ш		
					特記事項																										

キイトルーダは最大35サイクルまで。

インラインフィルター(0.2~5 µm)を使用すること。※カペシタビンは1日目の夕食後より服用開始。

	-								
カペシタビン減量基準	1日投与量								
ガベジダビン減重基準	初回投与量	減量段階1	減量段階2						
1.36㎡未満	2400mg/日	1800mg/日							
1.36㎡以上1.41㎡未満		Touting/ [1200mg/日						
1.41㎡以上1.51㎡未満	3000mg/日								
1.51㎡以上1.66㎡未満		2400mg/日							
1.66㎡以上1.81㎡未満	3600mg/日		1800mg/日						
1.81㎡以上1.96㎡未満	3000mg/ 🗖		Touting/ [
1.96㎡以上2.11㎡未満	4200mg/日	3000mg/日							
2.11㎡以上	4200ing/ 🗖		2400mg/日						

		L Tooting/ H										
	減量規定											
薬品名	エルプラット	カペシタビン										
標準投与量	130 mg∕ m i̇́											
1段階減量	100 mg∕ m i̇́	上記参照										
2段階減量	75 mg∕ m i̇́											
3段階減量	50 mg∕ m ²	中止(CCr≦30)										