

治療計画書

療法名 キイトルーダ(3週毎)+シスプラチン+RT 療法 1サイクル 3 週

疾患名 局所進行子宮頸癌

注射 内服 放射線	Rp	商品名	用量(／回)	投与ルート	投与時間	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
			用量(／日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
		照射部位	照射量																														
注射	1	生理食塩液	50 ml	点滴	15分																												
						●																											
注射	2	キイトルーダ	200 mg/body	点滴	30分																												
		生理食塩液	100 ml			●																											
注射	3	生理食塩液	50 ml	点滴	15分																												
						●																											
注射	4	ヴィーンF	500 ml	点滴	2時間																												
		硫酸Mg補正液20mEq/20ml	8 ml			●							●								●												
注射	5	ヴィーンF	500 ml	点滴	2時間																												
						●							●								●												
注射	6	グラニセトロン	3 mg	点滴	30分																												
		デキサート	9.9 mg			●							●								●												
注射	7	シスプラチン	40 mg/m ²	点滴	2時間以上																												
		生理食塩液	500 ml	全量500mlに調製		●							●								●												
注射	8	ヴィーンD	500 ml	点滴	2時間																												
		フロセミド	10 mg			●							●								●												
注射	9	ソリタT3	500 ml	点滴	2時間																												
						●							●								●												
注射	10																																
内服	1	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●						●	●						●	●												
内服	2																																
放射線			50 Gy	1.8-2Gy/day		●	●	●	●	●			●	●	●	●				●	●	●	●										
特記事項																																	
シスプラチンは最大5～6サイクルまで。 それ以降はキイトルーダ維持療法としてRp1～3を投与。(キイトルーダの投与量は、1～5サイクル目が200mg/3週毎、6～20サイクル目が400mg/6週毎で投与)																																	
減量規定																																	
薬品名	シスプラチン																																
標準投与量	40 mg/m ²																																
1段階減量	30 mg/m ²																																
2段階減量	中止																																
3段階減量																																	