

治療計画書

療法名FP(800+80)+キイトルーダ(3週毎)

療法1サイクル3週

疾患名根治切除不能な進行・再発の食道癌

注射	Rp	商品名	用量(/回)	投与ルート	投与時間	1						2					3							4							
内服			用量(/日)	用法	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
放射線		照射部位	照射量																												
注射	1	ヴィーンF	500 ml	点滴	2時間																										
		硫酸Mg補正液20mEq/20ml	8 ml			●																									
注射	2	生理食塩液	500 ml	点滴	2時間																										
						●																									
注射	3	キイトルーダ	200 mg/body	点滴	30分																										
		生理食塩液	100 ml			●																									
注射	4	生理食塩液	50 ml	点滴	10分																										
						●																									
注射	5	アロキシ	0.75 mg	点滴	30分																										
		デキサート	9.9 mg			●																									
		アロカリス	235 mg																												
注射	6	生理食塩液	50 ml	点滴	10分																										
						●																									
注射	7	シスプラチン	80 mg/m ²	点滴	2時間																										
		生理食塩液	500 ml	全量500mlに調製		●																									
注射	8	デキサート	6.6 mg	点滴(側管)	15分			●	●	●	●																				
		生理食塩液	100 ml																												
注射	9	ヴィーンF	500 ml	点滴(側管)	3時間			●	●	●	●																				
注射	10	5-FU	800 mg/m ²	点滴	24時間			●	●	●	●	●																			
		生理食塩液	500 ml																												
注射	11	フロセミド	10 mg	静注(側管)				●																							
注射	12	ヴィーンF	500 ml	点滴(側管)	2時間			●																							
						●																									
注射	13	生理食塩液	50 ml	点滴	15分						●																				
内服	1																														

特記事項

インラインフィルター(0.2~5μm)を使用すること。
キイトルーダ、5-FUは最大35サイクル。シスプラチンは最大6サイクル。

減量規定

薬品名	シスプラチン	5-FU	キイトルーダ		
標準投与量	80 mg/m ²	800 mg/m ²	200 mg		
1段階減量	60 mg/m ²	600 mg/m ²	中止		
2段階減量	40 mg/m ²	400 mg/m ²			
3段階減量					