

治療計画書

療法名 イミフィンジ+ゲムシタビン+シスプラチン(CCr40以上60未満) 療法 1サイクル 3 週

疾患名 膀胱がんにおける術前補助療法(4サイクルまで)

注射 内服 放射線	Rp	商品名	用量(/回) 用量(/日)	投与ルート 用法	投与時間	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
		照射部位	照射量																														
注射	1	生理食塩液	500 ml	点滴	2時間																												
						●							●																				
注射	2	イミフィンジ	※1500 mg/body	点滴	60分																												
		生理食塩液	100 ml			●																											
注射	3	生理食塩液	500 ml	点滴	2時間																												
						●							●																				
注射	4	グラニセトロン	3 mg	点滴	30分																												
		デキサート	13.2-16.5 mg			●							●																				
注射	5	ゲムシタビン	1000 mg/m <sup>2</sup>	点滴	30分																												
		生理食塩液	100 ml			●							●																				
注射	6	生理食塩液	50 ml	点滴	15分																												
						●							●																				
注射	7	シスプラチン	35 mg/m <sup>2</sup>	点滴	2時間																												
		生理食塩液	500 ml	全量500mlに調製		●							●																				
注射	8	ピカネイト	500 ml	点滴	2時間																												
		フロセミド	10 mg			●							●																				
注射	9	ヴィーンF	500 ml	点滴	2時間																												
						●							●																				
注射	10																																
内服	1	デカドロン	8 mg	1日1(-2)回朝(昼)食後		●	●	●	●	●			●	●	●	●	●																
内服	2																																
内服	3																																
内服	4																																
内服	5																																
内服	6																																
放射線			Gy																														
特記事項																																	
※体重30kg以下の場合、イミフィンジは20mg/kgで投与。 インラインフィルター(0.2又は0.22μm)を使用すること。																																	
減量規定																																	
薬品名																																	
標準投与量																																	
1段階減量																																	
2段階減量																																	
3段階減量																																	